題號:275

科目:熱力學(E)

題號: 275 共 [頁之第 全 頁

Please mark your answers clearly.

1. True or false? (5% each)

- A. The increase in the internal energy of a system is equal to the amount of energy added by heating the system minus the amount lost as a result of the work done by the system on its surroundings.
- B. Internal energy and entropy are intensive state variables.
- C. For a reversible adiabatic process in a simple system, the final temperature is a single-valued function of the final volume for a given initial state.
- D. As a system approaches absolute zero, all processes cease and the entropy of the system approaches a maximum value.
- E. The energy of an isolated system remains constant when a reversible process occurs in the system.

Please provide derivation and calculations for the following questions:

2. For the truncated viral equation of state:

$$\left(\frac{\partial \overline{U}}{\partial \overline{V}}\right)_{T,n} = \frac{RT^2}{\overline{V}^2} \frac{dB_2}{dT} = \frac{RT^2B'_2}{\overline{V}^2}$$

where R is the gas constant and \overline{V} is the molar volume. For argon gas at 298.15 K, B_2 is equal to -15.8 cm³/mol and B'_2 is equal to 0.25 cm³/mol·K. The molar constant-volume heat capacity of argon gas is nearly constant and equal to 3R/2. Find the internal energy, heat, and work for a reversible isothermal expansion of 1.0 mole of argon at 298.15 K from a volume of 2 L to a volume of 20 L. Calculate also these values assuming ideal gas behavior (30%).

- 3. If 2 moles of supercooled liquid water at -15°C freezes irreversibly at a constant pressure of 1 atm to ice, calculate the entropy change of the system, the surroundings, and the universe. Assume the surroundings remain at equilibrium at -15°C and the molar heat capacity of liquid water to be constant and equal to 76.1 J/mol·K and that of ice to be constant and equal to 37.15 J/mol·K (30%).
- 4. Find the expression for the partial molar Helmholtz energy of a one-component ideal gas as a function of pressure (15%).